



WHAT IS CLAIMED IS:

about 100,000 to about 500,000 IU/m^2 body surface/day.

comprising admin	A method for the potentiation of an immune response to an infectious scontinuation of treatment for the infection or prior to vaccination, istering to a subject a composition comprising IL-2 in an amount effective to enhancement without eliciting toxicity of Grade 1 or higher as defined by Organization.
2. Immunodeficiency	The method of claim 1, wherein the subject has Human Virus.
3. viral RNA.	The method of claim 2, wherein the subject has undetectable plasma
4.	The method of claim 3, wherein the subject has discontinued HAART.
5.	The method of claim 1, wherein the subject has HCV.
6. viral RNA levels.	The method of claim 5, wherein the subject has undetectable plasma
7. therapy or combina	The method of claim 6, wherein the subject has discontinued IFN- α tion IFN- α and ribavirin therapy.
8.	The method of claim 1, wherein the amount of IL-2 administered is

1
1
2
1
1
2
3
4
W ,
ĹÚ
47
1
3
<i>ý</i>

- 9. The method of claim 1, wherein the subject is immune reconstituted.
- 10. The method of claim 1, wherein the IL-2 composition further comprises a pharmaceutical carrier.
- 11. A method for treatment of hepatitis C virus comprising administering to a subject a composition comprising IL-2 in an amount effective to maintain immune enhancement without eliciting toxicity of Grade 1 or higher as defined by the World Health Organization.
- 12. The method of claim 11, wherein the IL-2 is administered in addition to IFN- α and ribavirin antiviral therapy.